

Probabilistic Method and Random Graphs

Lecture 8. The Method of Counting&Expectation

Xingwu Liu

Institute of Computing Technology, Chinese
Academy of Sciences, Beijing, China

¹The slides are mainly based on Chapter 6 of Probability and Computing.

Comments, questions, or suggestions?

Recap of Lecture 7

- Random graphs with fixed degree distribution
- Preferential attachment, rewired ring
- The existence of (sharp) threshold functions
 - Mostly on $\mathcal{G}_{n,p}$
 - Sharp threshold functions of connectivity: $\frac{\ln n}{n}$
 - That of major component existence: $\frac{1}{n}$
- A Hamiltonian cycle can be found efficiently w.h.p. in $\mathcal{G}_{n,p}$ with $p \geq 40 \frac{\ln n}{n}$

Probabilistic Method

-Elegance from graph theory

- A warm-up example:
 - n players against each other
 - “Top- k ” players get prize
- But, are you sure no **controversy** exists?
 - Controversy: prize-winners all defeated by a loser
- Unfortunately, when players are too many, controversy does exist w.h.p.



Proof (non-constructive)

- S : a k -subset of players
- A_S : no player beats all member of S
- Consider a random tournament


- $\Pr(A_S) = (1 - 2^{-k})^{n-k}$
- $\Pr(\text{no controversy}) = \Pr(\cup A_S)$
 $\leq \sum \Pr(A_S) = \binom{n}{k} (1 - 2^{-k})^{n-k}$
 $= o\left(\frac{1}{n}\right)$
- Find a controversial one? Just sampling

Cool?

A piece of cake in probabilistic method!

What is the Probabilistic Method?

- **Proving the existence** of an object satisfying a certain property **without constructing** it
- Underlying principle



- Pioneered by Erdős in 1940's
- Construct by flipping coins
- Naturally lead to (randomized) algorithms

Main Probabilistic Methods

- Counting argument
- First-moment method
- Second-moment method
 - Higher-moment method
- Lovasz local lemma

Algorithms

Counting Argument

- Construct a probability space and calculate the probability
- Algorithm design: sampling
- Application
 - Tournament
 - Ramsey number: an observation by S. Szalai, 1950's

Ramsey Number

- Given a complete graph $K(n)$, 2-color its edges
- Ramsey number $R(k, l)$
 - the smallest number n such that for every complete graph G with at least n vertices, any 2-coloring of G would either has a red $K(k)$ or a blue $K(l)$

There is a coloring without
red $K(k)$ and blue $K(l)$

$R(k, l)$

Every coloring has either
a red $K(k)$ or a blue $K(l)$

Number of vertices of the complete graph

Ramsey Number is Well defined

- Ramsey Theorem: $R(k, l)$ is finite for any k, l
 - Proved by F. Ramsey in 1930 and determined $R(3,3)$
 - Origin of Ramsey theory
 - The existence of rather large good substructure in a big structure
 - How much is $R(k, l)$?
- Upper bound: $R(k, l) \leq R(k - 1, l) + R(k, l - 1)$
 - Proved by P. Erdős and G. Szekeres in 1935
 - The 2nd cornerstone of Ramsey theory
 - By $R(k, 2) = R(2, k) = k$, $R(k, l) \leq \binom{k + l - 2}{k - 1}$
 - It implies $R(k, k) \leq 4^k$
 - Best: $k^{-c \frac{\ln k}{\ln \ln k}} 4^k$ by Conlon in 2009

Known bounding ranges

$r \backslash s$	1	2	3	4	5	6	7	8	9	10
1	1	1	1	1	1	1	1	1	1	1
2		2	3	4	5	6	7	8	9	10
3			6	9	14	18	23	28	36	40–42
4				18	25 ^[5]	36–41	49–61	59 ^[10] –84	73–115	92–149
5					43–48	58–87	80–143	101–216	133–316	149 ^[10] –442
6						102–165	115 ^[10] –298	134 ^[10] –495	183–780	204–1171
7							205–540	217–1031	252–1713	292–2826
8								282–1870	329–3583	343–6090
9									565–6588	581–12677
10										798–23556

Vigleik Angeltveit; Brendan McKay (2017). “ $R(5,5) \leq 48$ ”. [arXiv](#)


Proof of the upper bound

- 2-color the complete graph on $R(k - 1, l) + R(k, l - 1)$ vertices
- Pick a vertex u
- Define subgraphs G_r and G_b : $\forall v$,
 - $v \in \begin{cases} G_r & \text{if } (u, v) \text{ is red} \\ G_b & \text{if } (u, v) \text{ is blue} \end{cases}$
- Either $|G_r| \geq R(k - 1, l)$ or $|G_b| \geq R(k, l - 1)$
 - Do case-by-case analysis

Example: Ramsey Number $R(3,3)$

$$R(3,3) \leq 6$$

Actually, $R(3,3) > 5$



What is $R(k, k)$?


- $R(k, k) > 2^{k/2} = \sqrt{2}^k$ (Erdős, 1947)
 - Best: $[1 + o(1)] \frac{k}{e} \sqrt{2}^{1+k}$ by Spencer in 1975
- For any complete graph with at most $2^{k/2}$ vertices, there is a 2-coloring without red $K(k)$ and blue $K(k)$
- Prove by the **probabilistic method**

Proof

- Consider a graph G on n vertices, and 2-color each edge randomly
 - Uniform distribution on all 2-coloring
- A_S : the subgraph on S is monochromatic
 - S is a k -subset of the vertices
- $\Pr(A_S) = 2^{1-\binom{k}{2}}$
- $\Pr(\cup_S A_S) \leq \binom{n}{k} 2^{1-\binom{k}{2}} < \frac{2^{1+\frac{k}{2}} n^k}{k! \frac{k^2}{2}} < 1$ (If $n = \lfloor 2^{k/2} \rfloor$)
- There is a 2-coloring of the edges such that $K(\lfloor 2^{k/2} \rfloor)$ has no monochromatic k subgraph
 - Together with the definition of $R(k, k)$, $R(k, k) > 2^{k/2}$

Randomized Algorithms

- But how to find a good coloring? By sampling!
- General approach



- Prerequisites
 - Efficient sampling
 - Small probability of failure
 - Efficient verification (Las Vegas only)

First-Moment method

- Use the expectation in probabilistic reasoning
- Two types of first-moment method

- Expectation argument

$$\Pr(X \geq \mathbb{E}[X]) > 0, \Pr(X \leq \mathbb{E}[X]) > 0$$

- Markov's inequality for non-negative X

- $\Pr(X \geq a) \leq \frac{\mathbb{E}[X]}{a}$

- When X is integer-valued,

$$\Pr(X \neq 0) = \Pr(X > 0) = \Pr(X \geq 1) \leq \mathbb{E}[X]$$

First-Moment argument

- 3-CNF Boolean formula
 - $(x_1 \vee x_2 \vee x_3) \wedge \dots \wedge (\overline{x_1} \vee \overline{x_3} \vee x_4)$
- For such a formula, at most how many clauses can be satisfied simultaneously?
 - **MAX-3SAT** is NP-hard
- Theorem: there is a truth assignment which satisfies $\geq \frac{7}{8}$ -fraction of the clauses

Proof

- Randomly assign truth values to each variable
- Define r.v. X_i indicating whether clause i is true
- $\mathbb{E}[X_i] = \frac{7}{8} \Rightarrow \mathbb{E}[\sum_{i=1}^n X_i] = \frac{7}{8}n$
- Remark: probability of sampling a good truth assignment $\geq \frac{1}{n+1}$, leading to an efficient alg.
 - Optimum, since impossible to get a $(\frac{7}{8} + \varepsilon)$ -approx.
 - J. Hastad. Some optimal inapproximability results. STOC 1997

Proof of $\Pr\left(\sum X_i \geq \frac{7}{8}n\right) \geq \frac{1}{n+1}$

- Let $X = \sum X_i$ and $p = \Pr\left(X \geq \frac{7}{8}n\right)$

- $\frac{7}{8}n = \mathbb{E}[X]$

$$= \sum_{i < \frac{7}{8}n} i * \Pr(X = i) + \sum_{i \geq \frac{7}{8}n} i * \Pr(X = i)$$

$$\leq \left(\frac{7}{8}n - \frac{1}{8}\right) (1 - p) + np$$

$$= \frac{7}{8}n - \frac{1}{8} + \frac{n+1}{8}p$$

Expectation argument

- Turán Theorem
 - Any graph $G=(V,E)$ contains an independent set of size at least $\frac{|V|}{D+1}$, where $D = \frac{2|E|}{|V|}$
- Proof: Consider the following **random process** for constructing an independent set S :
 - Initialize S to be the empty set
 - For each vertex u in V in random order,
 - if no neighbors of u are in S , add u to S
 - Return S

Proof (Continued)

- S is an independent set
- Vertex u is selected with probability $\geq \frac{1}{d(u)+1}$
 - See the next slide
- So, $\mathbb{E}[|S|] \geq \sum \frac{1}{d(u)+1} \geq \frac{|V|}{D+1}$ due to convexity
- **Remark:** probability of sampling a good independent set is $\geq \frac{1}{2D|V|^2}$

$$\text{Proof: } \Pr(u \text{ is selected}) \geq \frac{1}{d(u)+1}$$

- u is selected if and only if A occurs
 - A : when sampling first occurs in the neighborhood of u , u rather than its neighbors is sampled
 - Neighborhood: u and its then-valid neighbors
 - Denote the neighborhood by N , and the number of then-valid neighbors by x . Note that $x \leq d(u)$
- $\Pr(A) = \Pr(u \text{ is chosen} \mid \text{sampling occurs in } N)$
$$= \frac{1}{x+1} \geq \frac{1}{d(u)+1}$$

References

- <http://www.cse.buffalo.edu/~hungngo/classes/2011/Spring-694/lectures/sm.pdf>
- <http://www.cse.cuhk.edu.hk/~chi/csc5160-2007/notes/Probabilistic.ppt>
- Erdős. Graph theory and probability I. 1959
- Erdős. Graph theory and probability II. 1961
- Alon&Krivelevich. Extremal and Probabilistic Combinatorics. 2006